



BioBacta



Journal of Bioscience and Applied Research

www.jbaar.org



Isolation of *Aspergillus* section *Flavi* and determination of aflatoxins in Bambara groundnut sold in Cotonou main markets (Benin)

Yann Christie Sissinto ADJOVI¹, *, Prudence AGNANDJI², Lucie AYI-FANOU², and Ambaliou SANNI²

1. Unit of Biochemistry and Molecular Biology, National University of Agriculture, Benin

-Ecole des Sciences et Techniques de Conservation et de Transformation des Produits Agricoles / Université Nationale d'Agriculture, Mail: yann.adjovi6@gmail.com **HYKK3RBY5N**

Adress : Unité de Biochimie et de Biologie Moléculaire, 04 P.O. Box 0320, Cotonou, Benin

2. Unit of Biochemistry and Molecular Biology /University of Abomey-Calavi

Adress: Unité de Biochimie et de Biologie Moléculaire, 04 P.O. Box 0320, Cotonou, Benin

*Corresponding author: yann.adjovi6@gmail.com

Received: November 1, 2019; Accepted: December 2, 2019

DOI :10.21608/jbaar.2019.115437

Abstract

Legumes such as cowpea (*Vigna unguiculata*) and Bambara groundnut (*Vigna subterranean*) are the most sought in Benin and in the West Africa sub-region. These legumes are an important protein source because of their high content of essential amino acids. But, like many dried agricultural and food products, they are subject to contamination by molds and mycotoxins during storage.

In order to assess the contamination of Bambara groundnuts, by the most dangerous and widespread mycotoxins in Africa (aflatoxins), especially in Benin, this study was conducted in Cotonou (Benin) main markets. Thirty-five (35) samples of Bambara groundnuts were collected from six (6) different markets in Cotonou. The fungi are isolated and identified morphologically on different media after direct culture. The species of *Aspergillus* section *Flavi* aflatoxigenic potential are evaluated and the contamination of samples by aflatoxins was determined by Thin Layer Chromatography (TLC).

The results of this study showed that the average water content of Bambara groundnuts samples ranged from 4.85 to 9.35%. The main species of mold isolated are *Mucor spp.*, *Rhizopus spp.*, *Alternaria sp.*, and *Aspergillus spp.* Forty-six (46) strains of *Aspergillus* section *Flavi* were isolated on Potatoes Dextrose Agar (PDA). All of them have been identified as aflatoxigenic strains and were divided into different groups according to their morphological appearance and secondary metabolites. Twenty-two samples (62.85%) of Bambara groundnuts were contaminated by aflatoxins.

This study showed that Bambara groundnuts which sold in the main markets of Cotonou constitute a health risk for populations.

Keywords: Bambara groundnut, *Vigna subterranean*, *Aspergillus flavus*, aflatoxins.

1. INTRODUCTION

To have a balanced diet, humans need nourishing food that contains essential nutrients. The nutrients needed by humans are proteins that span structure (growth, tissue repairs), while fats and carbohydrates generally provide the energy needs (Lagnika *et al.*, 2017; Mathieu-Daude *et al.*, 2001). Foods rich in these elements are cereals and legumes such as cowpea and Bambara groundnut that are consumed in Africa.

Although the environment has all elements necessary for organism survival, various health problems jeopardize people's health, especially in developing countries such as Benin (Zinzindohoue, 2012). Indeed foodstuffs (cereals, tubers, legumes, vegetables, etc.) (Dudler, 2005) are subject to various contaminations among which are those caused by mycotoxins. These are produced by toxigenic molds of the genus *Aspergillus* (*A.*), *Penicillium* (*P.*) and *Fusarium* (*F.*) (Zinedine and Idrissi, 2007). Aflatoxin B1 is the most dangerous because of its toxicity. Aflatoxins pose serious health hazards to humans and domestic animals because they frequently contaminate agricultural commodities. In 1993, Bayman *et al* showed that aflatoxin produced by *Aspergillus flavus* (*A. flavus*), *Aspergillus parasiticus* (*A. parasiticus*) and *Aspergillus nomius* (*A. nomius*) are highly teratogenic, mutagenic and carcinogenic in animals and in human. In 2007, Njapau *et al.* reported the death of about one hundred and twenty-five people after consuming contaminated corn with aflatoxins in Kenya. Aflatoxins produced by *A. flavus* and *A. parasiticus* contaminate foods such as rice, corn, peanuts, spices etc (Cho *et al.*, 2008 ;

Gnonlonfin *et al.* 2012; Jalili and Jinap, 2012; Serra *et al.*, 2002).


Recognition of the need to control aflatoxin contamination of food and feed grains has elicited various approaches from researchers to eliminate this toxin from maize and other susceptible crops. The approach to enhance host resistance through conventional means has gained renewed attention following the discovery of natural resistance to *A. flavus* infection and aflatoxin production in many crops like maize (Brown *et al.*, 2001) cassava (Adjovi *et al.*, 2015; Adjovi *et al.*, 2014; Gnonlonfin *et al.*, 2008) and cowpea (Houssou *et al.*, 2009). In this last product, Houssou *et al.* have proposed the presence of protein which confers this property to cowpea. This study focused on the research of aflatoxins and *Aspergillus* section *Flavi* on Bambara groundnut (*Vigna subterranean*), a legume of the same genus with cowpea. The aim is to investigate if Bambara groundnuts stored and sold in Benin are contaminated by *Aspergillus* section *Flavi* aflatoxinogenic's strains and aflatoxins or have a resistance property as cowpea.

2. MATERIALS AND METHODS

Sampling

Thirty-five (35) samples of Bambara groundnut (*V. subterranea*) were collected from six (6) main markets in Cotonou (Gbegamey, Kindonou, Dantokpa, Fijrossè, Akpakpa, and Ganhi) (figure 1). Sampling was done from different traders in order to have heterogeneous samples. Samples were transported to the laboratory and stored at 4°C for different analysis.

Figure 1: Sampling



Water content

Water content is determined by AOAC (Association of Official Analytical Chemist) method 934.01

AOAC. 5 g of Bambara groundnut (*V. subterranea*) are ground in a blender and dehydrated at 70 ° C.

After 20 hours, the mass of Bambara groundnut (*V. subterranea*) is measured. Then, moisture is determined in the percentage of wet mass.

Mycological analysis

The direct plating method described by Ravi Kiran et al. (2005) was used with some modifications. Ten seeds of each sample of Bambara nuts are surface sterilized with 10% bleach (potassium hypochlorite solution) for 1 min and rinsed twice with sterile distilled water. Three seeds of Bambara nuts were placed on Dichloran Chloramphenicol 18% Agar (DG 18) (Oxoid Ltd, Hampshire, UK) in Petri dishes and incubated at 25 °C for 7 days. Triplicate of each sample analysis are done. After this, each Fungi colony are transplanted on the center of Potatoes Dextrose Agar (PDA) and incubated at 30°C for 7 days.

Aspergillus section Flavi identification

The total number of fungal isolated from each product was recorded. Identification of *Aspergillus section Flavi* was firstly based on morphology. Macroscopic and microscopic features of each colony were used. A loop full of pure isolate cultured on potatoes dextrose agar (PDA) was sub-cultured on malt extract agar (MEA), Yeast Extract Agar (YES) and *Aspergillus Flavus Parasiticus* Agar (AFPA) (Oxoid Ltd., Hampshire, UK) for identification. The MEA and YES plates were incubated at 25°C for 7 days. The identification of fungi was based on gross morphology and microscopic features such as spores and fruiting structures using Pitt and Hocking (2009) descriptions.

Detection of aflatoxigenic strains

- Extraction

Aflatoxins were extracted from 7 days cultures of *Aspergillus Section Flavi* strains on malt extract agar medium. Each culture are extracted with methanol/water (80/20, V / V). The extracts are dry

at 60°C. The dry extract obtained is taken up in methanol 400µl.

- Aflatoxins detection

Aflatoxins spot were detected by Thin Layer Chromatography. 10 µL of extracts and standard of Aflatoxins (Sigma Aldrich) were deposited on the fluorescent chromatographic plate (1.05553. DC-Alufohlen Kieselgel 60). Migration solvent is ether / methanol / water (96: 3: 1, v / v / v). The plates placed in the solvent system were dried and observed at UV 365nm.

Detection of Aflatoxins in Bambara groundnuts

Aflatoxins were extracted and analyzed according to the method of Bankole and Mabekoje (2004). 5 g of Bambara were ground, mixed with twenty-five (25) ml of methanol / water (85: 15v / v) and stirred for three (3) minutes. The mixture was filtered through Whatman filter paper N° 1. Twenty (20) ml of the filtrate were mixed with twenty (20) ml of sodium chloride (10% NaCl) and shaken well. The mixture was transferred to a separatory funnel and with 10 ml of hexane. The hexane layer was discarded and the aqueous phase was recovered. 25 ml of chloroform was added to the aqueous phase. The Chloroform phase was rejected. The extract obtained was filtered through Whatman N°1 containing 2.5 g of anhydrous sodium thiosulfate. The filtrate obtained was evaporated with a rotary evaporator (RE 300). The eluate was dried down at 60°C and then reconstituted in 50 µL of methanol and stored at 4°C until analysis. Analysis of extracts obtained from the Bambara groundnut (*V. subterranean*) is done as previously described by Thin layer chromatography.

Statistical analysis

Data were analyzed with ANOVA for multiple comparisons among means. Correlation and regression coefficients were calculated using SPSS for Windows version 18.0 (SPSS, Chicago, IL).

3. RESULTS

Water content

Average water content is recorded in the table below. These averages vary from 4.85% to 9.35% (Table 1).

Table 1: Average of sample water content

Sites	Akpakpa	Gbégamey	Dantokpa	Ganhi	Fidjrossè	Kindonou
Moy %	5,85	5,85	6,40	8,45	9,35	4,85
Ecart types	0,003	0,016	00	0,002	0,002	0,004

Mycological analysis

Bambara groundnut or *V. subterranea* are more contaminated by species of *Aspergillus spp*, *Rhizopus spp*, *Mucor spp*, and *Alternaria sp*. The prevalence of each genus are respectively:

In *Aspergillus* species, two main dominant species contaminate Bambara samples, *A. niger* (incidence 67.38) followed by *A. flavus* (27.22). The incidence of this last species varies between 5.28% in Kindonou's samples and 31.87% in Fidjrosse's samples.

Samples from Fidjrosse's market were the most contaminated with *A. flavus*. The ANOVA Turkey made an error rate of 5% has allowed us to find a significant difference between the samples from Fidjrosse and those from Dantokpa, Ganhi, Gbégamey, Akpakpa, and Kindonou. Samples from Kindonou seem the least contaminated.

Forty-six (46) strains isolated have been morphologically identified as strains of *Aspergillus* section *Flavi* and two as *Aspergillus* section *Tamaritii*. The repartition of *Aspergillus* section *Flavi* strains by sight is as follows: Dantokpa (12), Akpakpa (10),

Ganhi (8), Fidjrosse (7), Gbégamey (6) and Kindonou (5).

Detection of *Aspergillus* section *Flavi* isolated nature

The analysis of media extract revealed that all of the 46 strains isolated are aflatoxinogenic. The aflatoxinogenics strains were classified according to the different types of aflatoxin they produce in Table 2.

The culture on AFPA media show that all strains are aspergilliacid producers (Table 2).

These results show that the Bambara groundnut (*V. subterranea*) sold in the city of Cotonou is contaminated with strains of aflatoxinogenic *Flavi* section. Samples from Akpakpa markets Dantokpa and Fidjrosse are more contaminated than from Ganhi, Gbégamey and Kindonou.

Aflatoxins detection in Bambara groundnuts samples

The results of the qualitative analysis of collected samples carried out show that 22 of the 35 samples are infected by aflatoxins. The results are shown in Table 3 following.

Table 2: Type patterns of *Aspergillus* section *Flavi* strains based on aflatoxins, aspergillitic acid and sclerotia production

Types	Sclerotia	Aflatoxins	Aspergillitic acid	Related species	Number of strains
I	None	AFB1	+	<i>A.flavus</i>	13
			+	<i>A.parasiticus</i> , <i>A.bombicys</i> , <i>A.novoparasiticus</i> , <i>A.arachidichola</i> , <i>A.pseudocaelatus</i>	
II	None	AFB1, AFB2, AFG1, AFG2			4
III	Large and abundant	AFB1	+	<i>A.togoensis</i>	5
			+	<i>A.flavus</i> , <i>A.pseudonomius</i> , <i>A.pseudotamarii</i>	
IV	Large and abundant	AFB1, AFB2	+	<i>A.transmontanensis</i> , <i>A.sergii</i> , <i>A.toxicarius</i>	8
V	Large and little	AFB1, AFB2, AFG1, AFG2			
VI	Small and little	AFB1	+	<i>flavus clade</i>	1
			+	<i>A.minisclerotigenes</i> , <i>A.parvisclerotigenus</i>	
VII	Small and little	AFB1, AFB2, AFG1, AFG2			10

(Ito et al., 2001; White and Hill, 1943; Saito and Tsuruta, 1993; (Frisvad, Skouboe and Samson, 2005) (M. B. Pildain et al., 2008; Rank et al., 2011; Varga, Frisvad and R. A. Samson, 2011; Soares et al., 2012; Adjovi et al., 2014)

Table 3: Aflatoxins contaminant in Bambara groundnuts (*V. subterranea*) in Cotonou sample


Sites	Samples	AFB1	AFB2	AFG1	AFG2
Akpakpa	E ₁ A	-	-	-	-
	E ₂ A	-	-	-	-
	E ₃ A	-	-	-	-
	E ₄ A	+	+	+	-
	E ₅ A	+	+	+	-
Fidjrossè	E ₁ F	+	-	+	-
	E ₂ F	+	+	+	-
	E ₃ F	+	-	+	-
	E ₄ F	+	-	+	-
	E ₅ F	+	+	+	-
Gbégamey	E ₆ F	+	-	-	-
	E ₁ G	+	-	-	-
	E ₂ G	-	-	-	-
	E ₃ G	+	+	-	-
	E ₄ G	-	-	-	-
Kindonou	E ₅ G	+	-	-	-
	E ₆ G	-	-	-	-
	E ₁ K	+	+	+	-
	E ₂ K	+	-	-	+
	E ₃ K	+	-	-	-

4. DISCUSSION

Usually, foods are contaminated by different microorganisms like molds (Marzia *et al.*, 2014). This contamination is often favored by the high water activity of the substrate, but also by other factors such as the nature of the substrate and temperature (Yu and Ehrlich, 2011). However, it has been proven that foods with low water content are also contaminated by mold (Schatzki *et al.*, 2002). These findings demonstrate that *A. flavus* infection is not only due to water activity but also of inappropriate post-harvest handling methods used by farmers and other value

chain actors. This is the case of the seeds of Bambara groundnut (*V. subterranea*) that have low water activity (4.85% to 9.35%) but are nevertheless highly contaminated by fungi. Similar results were found by Schatzki *et al.* (2002) and Houssou *et al.* (2009) who worked respectively on cowpea and groundnut (legume) whose water content ranged respectively from 6, 27-8% and 8, 8-12.3% and have high fungi contamination. Figure 2 shows the correlation between Bambara groundnut water content and *Aspergillus* section *Flavi* contamination.

Figure 2: Correlation between Bambara groundnut contamination by *A. flavus* strains and the water content



In this study, Bambara groundnut collected in Cotonou's main market is mainly contaminated by genus *Mucor*, *Rhizopus*, *Aternaria* and *Aspergillus*. Usually, Bambara groundnut (*V. subterranea*) is harvested, sun-dried and sold without treatment, which can promote the contamination by soil molds and storage conditions contribute to storage's fungi development. Hot and humid climates, prevalent in the tropics promote fungi that infect the crop in the field and/or immediately after harvest during storage (IARC, 2002). Also, Houssou *et al.* (2009) reported that among the factors contributing to contamination by molds of the genus *Aspergillus*, storage plays an

important role. This observation is confirmed by (Seetha *et al.*, 2018) who have noted the contamination of fresh Bambara groundnut by fungi like *Aspergillus flavus* which increase during storage. Virtually all of our samples are contaminated by molds of the genus *Aspergillus* with an average of CFU up to 94.68% and 50 strains isolated. Among this genus, the species *A. niger* had the most prevalence (67.38%) followed by the section *Flavi* (prevalence of 27.22%) with 46 strains identified and the last *Aspergillus* section *Tamaritii* with 2 strains. The prevalence of *Aspergillus* section *Flavi* in this work is less than in a study in Durban which shows

the high presence of *Aspergillus* section *Flavi* in 64.7% of the Bambara groundnut sample (Olagunju *et al.*, 2018).

The identification of species and aflatoxigenic nature of *Aspergillus* section *Flavi* isolated on Bambara groundnuts in Benin, reveal that most of them produce sclerotia and all of the 46 isolates, produce aflatoxins (Table 2). Strains have been regrouped into seven types based on aflatoxins type produced, sclerotia size and aspergillitic acid production. It's known that all of *Aspergillus* of section *Flavi* is not *Aspergillus flavus* var. *flavus* which is characterized by the production of aflatoxins B1 only (Varga *et al.*, 2011). Many authors have shown evidence that *A. flavus sensu lato* may consist of several species (Geiser *et al.*, 1998); (B. Pildain *et al.*, 2008); (Soares *et al.*, 2012)). Diversity in the population of *Aspergillus* section *Flavi* exists with respect to their size of sclerotia (Kiewnick *et al.*, 2005). S-strain (those producing small sclerotia) produce high amounts of aflatoxins and L-strains with large sclerotia (Cotty and Cardwell, 1999; Kiewnick *et al.*, 2005). Chemotaxonomy permits also identification of *Aspergillus* section *Flavi* species. The group in section *Flavi* is composed of (i) an *A. flavus* clade containing *A. flavus*, *A. parvisclerotigenus* and *A. minisclerotigenes*, (ii) an *A. parasiticus* clade containing *A. parasiticus*, *A. arachidicola*, *A. transmontanensis* and *A. sergii* and (iii) *A. mottae* as a sibling to the clade containing both *A. parasiticus* and *A. flavus* (Soares *et al.*, 2012). In this work, types called type I, III, IV, VI and VII can be identified as *Aspergillus flavus* clade *flavus*, with type III as *A. togoensis* causes of the similitudes like described by (Rank *et al.*, 2011) who produce AFB1 only and large sclerotia; type II and V can be related to *A. parasiticus* clade.

The presence of aflatoxins have been revealed in Bambara groundnuts produced and sold in Cotonou mains markets. In this work, 22 of the 35 samples collected (62.86%) were infected by aflatoxins. Maringe *et al.* (2017) have found 3 out of 89 (3.4%) Bambara nut samples contaminated by aflatoxins in

South Africa which is very small compared to our results. Another study of (Seetha *et al.*, 2018) shows the contamination of fresh and stored Bambara groundnut by aflatoxins in central Tanzania.

This result confirms and shows that Bambara groundnut (*V. subterranean*), does not contain aflatoxins inhibitory substances and appears to be a real danger in Benin such as peanuts, corn, and cereal as favorable substrates for aflatoxins production.

CONCLUSION

At the end of our study on the evaluation of the contamination of the Bambara groundnut (*Vigna subterranae*) in Benin, it appears that the Bambara groundnut (*Vigna subterranae*) produced and sold in Benin is subject to contamination by molds of the genus *Mucor*, *Rhizopus*, *Alternaria*, and *Aspergillus*. Many strains of *Aspergillus* section *Flavi* (46) of different clade isolated have been identified as aflatoxinogenic. 62.86% of the collected samples were found contaminated with different types of aflatoxins.

This study is another work to reveal the importance of the health hazard posed by the consumption of food contaminated by mycotoxins in Benin.

ACKNOWLEDGEMENTS

We thank Maciana TANG for providing language help.

5. REFERENCES

- Adjovi, Y.C.S., Gnonlonfin, B.J.G., Bailly, S., Bailly, J-D., Tadrict, S., Puel, O., Oswald, P.O.I., Sanni, A. (2015) 'Occurrence of mycotoxins in cassava (*Manihot esculenta* Crantz) and its products'. International Journal of Food Safety, Nutrition and Public Health 5 (3/4), pp. 217-246.
- Adjovi, Y. C.S., Bailly, S., Gnonlonfin, B. J.G., Tadrict, S., Querin, A., Sanni, A., Oswald, I. P., Puel, O., Bailly, J. D. (2014) 'Analysis of the contrast between natural occurrence of toxigenic *Aspergilli* of the *Flavi* section and aflatoxin B1 in cassava', Food Microbiology, 38, pp. 151–159.

Brown, R. L., Chen, Z., Menkir, A., Cleveland, T. E., Cardwell, K., Kling, J. and White, D. G. (2001) Resistance to Aflatoxin Accumulation in Kernels of Maize Inbreds Selected for Ear Rot Resistance in West and Central Africa. *Journal of Food Protection* : March 2001, Vol. 64, No. 3, pp. 396-400.

Cho, S. H., Lee, C. H., Jang, M. R. (2008). “Aflatoxins contamination in spices and processed spice products commercialized in Korea,” *Food Chemistry*, 1. 107, pp. 1283–1288.

Cotty, P. J. and Cardwell, K. F. (1999) ‘Divergence of West African and North American Communities of *Aspergillus Section Flavi*’, *Society*, 65(5), pp. 2264–2266.

Dudler, Vincent. 2005. ordonnance sur les denrées alimentaires et les objets usuels (RS 817.02) Colloque IUMSP4.12.2012.

FAO (2004). Food and Agriculture Organization, Worldwide regulations for mycotoxins in food and feed in 2003. FAO, Food and Nutrition Paper N° 81. Rome, Italy.

FAO (2001). Manual on the application of HACCP system in mycotoxin prevention and control. Food and Agriculture Organisation of the United Nation, Rome.

Frisvad, J. C., Skouboe, P. and Samson, R. (2005) ‘Taxonomic comparison of three different groups of aflatoxin producers and a new efficient producer of aflatoxin B1, sterigmatocystin and 3-O-methylsterigmatocystin, *Aspergillus rambellii* sp. nov.’, *Systematic and Applied Microbiology*, 28(5), pp. 442–453.

Geiser, D. M., Pitt, J. I. and Taylor, J. W. (1998) ‘Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*’, *Proceedings of the National Academy of Sciences of the United States of America*, 95(1), pp. 388–93.

Gnonlonfin, G. J. B., Hell, K., Fandohan, P. and Siame, A. B. (2008) ‘Mycoflora and natural occurrence of aflatoxins and fumonisin B1 in cassava and yam chips from Benin, West Africa.’, *International journal of food microbiology*, 122(1–2), pp. 140–147.

Houssou, P., Ahohuendo, B. C., Fandohan, P., Kpodo, K., Hounhouigan, D. J. and Jakobsen, M. (2009) ‘Natural infection of cowpea (*Vigna unguiculata* (L.) Walp.) by toxigenic fungi and mycotoxin contamination in Benin, West Africa’, *Journal of Stored Products Research*. Elsevier Ltd, 45(1), pp. 40–44.

IARC, International Agency for Research on Cancer (2002). Monograph on the evaluation of carcinogenic risk to humans, vol. 82. World Health Organization, IARC, Lyon, France.

Ito, Y., Peterson, S. W., Wicklow, D. T. and Goto, T. (2001). *Aspergillus pseudo-tamarii* a new aflatoxin producing species in *Aspergillus* section *Flavi*. *Mycol. Res.* 105, pp. 233–239.

Jalili, M. and Jinap, S. 2012. “Natural occurrence of aflatoxins and ochratoxin A in commercial dried chili”, *Food Control*, 24, pp. 160–164.

Kiewnick, S. et al. (2005) ‘Tropentag 2005 Stuttgart-Hohenheim, October 11-13, 2005’, *Biological Control*, pp. 13–16.

Lagnika, C., Adjovi, Y. C. S., Lagnika, L., Gogohounga, Fructueux O., Do-Sacramento, O., Koulony, R. K., Sanni, A. (2017) ‘Effect of Combining Ultrasound and Mild Heat Treatment on Physicochemical, Nutritional Quality and Microbiological Properties of Pineapple Juice’, *Food and Nutrition Sciences*, 08(02), pp. 227–241.

Olagunju, O., Mchunu, N., Durand, N., Alter, P., Montet, D., Ijabadeniyi, O. (2018) ‘Effect of milling, fermentation or roasting on water

activity, fungal growth, and aflatoxin contamination of Bambara groundnut (*Vigna subterranea* (L.) Verdc)', *LWT*, 98, pp. 533–539.

Pildain, M.B., Frisvad, J.C., Vaamonde, G., Cabral, D., Varga, J., and Samson, R.A. (2008). Two novel aflatoxin-producing *Aspergillus* species from Argentinean peanuts. *International Journal of systematic and evolutionary microbiology* 58, pp. 725–735.

Rank, C., Nielsen, K. F., Larsen, T. O., Varga, J., Samson, R. A., Frisvad, J. C. (2011) 'Distribution of sterigmatocystin in filamentous fungi', *Fungal Biology*. Elsevier Ltd, 115(4–5), pp. 406–420.

Ravi Kiran, D., Narayana, K.J.P., & Vijayalakshmi, M. (2005). Aflatoxin B1 production in chillies (*Capsicum annum* L.) kept in cold stores. *African Journal of Biotechnology*, 4, pp. 791–795.

Saito, M. and Tsuruta, O. (1993). A new variety of *Aspergillus flavus* from tropical soil in Thailand and its aflatoxin productivity. *Proc. Japanese Associ. Mycotoxicol.* 37, pp.31–36.

Schatzki T. F. and Haddon W.F., (2002). 'Rapid, Non-destructive Selection of Peanuts for High Aflatoxin Content by Soaking and Tandem Mass Spectrometry'. *Journal of Agricultural and Food Chemistry.*, 50 (10), pp. 3062-3069.

Seetha, A., Monyo, E., W. Tsusaka, T., W. Msere, H., Madinda, F., Chilunjika, T., Sichone, E., Mbughi, D., Chilima, B., Matumba, L. (2018) 'Aflatoxin-lysine adducts in blood serum of the Malawian rural population and aflatoxin contamination in foods (groundnuts, maize) in the corresponding areas', *Mycotoxin Research*. 33 (4), pp. 323–331.

Serra I, Yamamoto M, Calvo A, Cavada, G., Báez, S., Endoh, K., Watanabe, H., Tajima, K. (2002) Association of chili pepper consumption, low socioeconomic status and longstanding

gallstones with gallbladder cancer in a Chilean population. *Int J Cancer*, 102, pp. 407-411.

Soares, C., Rodrigues, P., Peterson, S., Lima, N., Venâncio, A. (2012) 'Three new species of *Aspergillus* section *Flavi* isolated from almonds and maize in Portugal.', *Mycologia*, 104(3), pp. 682–97.

Varga, J., Frisvad, J. C. and Samson, R. (2011) 'Two new aflatoxin producing species, and an overview of *Aspergillus* section *Flavi*.' , *Studies in mycology*, 69(1), pp. 57–80.

White, E. C. and Hill, J. H. (1943) 'Studies on Antibacterial Products Formed by Molds : I. *Aspergill*ic Acid, a Product of a Strain of *Aspergillus flavus*', *Journal of Bacteriology*, 45(5), pp. 433–443.

Yu, J. and Ehrlich, K. C. (2011) 'Aflatoxin Biosynthetic Pathway and Pathway Genes'. *Aflatoxins – Biochemistry and Molecular Biology*. 10.5772/23034.

Zinedine, A. and Idrissi, L. (2007) 'Présence et réglementation des mycotoxines dans les aliments au Maroc : Situation actuelle et perspectives', pp. 10–18.