



Journal of Bioscience and Applied
Research

JBAAR

WWW.JBAAR.ORG



***Moringa oleifera* leaf Extract: A Potent Ameliorator of Cyclophosphamide Induced liver Toxicity in Rat Model**

Ahmed Abdulzahra Habeeb

Mayssan Health Directorate Higher Institute of Health, Iraq

DOI: [10.21608/JBAAR.2018.129732](https://doi.org/10.21608/JBAAR.2018.129732)

Abstract

Moringa oleifera leaf is pharmacologically active with documented antioxidant activity. In the current study protective effect of ethanol extract of *Moringa oleifera* leaf extract (MLE) was investigated in rats against cyclophosphamide (CYP) induced liver injuries. Twenty-eight Wistar albino rats were divided into four groups, as follows: 1) control group - received vehicle used for MLE and CYP for 14 days; 2) MLE group - rats were administered orally at a dose 200.0 mg kg⁻¹ b.wt. for 14 consecutive days; 3) CYP group - cyclophosphamide at a dose of 150 mg/kg was given through i.p. to rats as a single dose at day 7; 4) MLE + CYP group – MLE was given for 14 days plus a single dose of CYP was given on hour after MLE administration. Catalase (CAT), glutathione (GSH), the level of lipid peroxidation thiobarbituric acid-reactive substances (TBARS), DNA, and RNA concentration were analyzed in liver tissue. Also, serum total protein, albumin, cholesterol and triglycerides, aspartate aminotransferase (AST) and alanine aminotransferase (ALT), alkaline phosphatase (ALP), lactic dehydrogenase (LDH) values were evaluated. AST, ALT, ALP, LDH, triglycerides, and total cholesterol in serum were found to be significantly ($P < 0.05$) higher in the CYP group, compared to the control group, while protein and albumin were decreased. Compared with the control group, significantly high levels of liver TBARS and the low antioxidant defenses, like free radical scavenging enzyme viz., catalase activity as well as GSH concentration in CYP-treated group. In rats supplemented with MLE as well as treated with CYP, hepatic specific marker enzymes were restored to normalcy which otherwise was lowered in the CYP-treated rats. In conclusion, MLE exhibited antioxidant activity by the presence of free-radical quenching constituents.

Key Words: cyclophosphamide; *Moringa oleifera* leaf extract; Liver; Rat; Oxidative stress

Received: December 20, 2017. Accepted: February 27, 2018. Published: March 3, 2018

1. Introduction

Free radicals are the major by-products produced by the cells of aerobic organisms and can start the autocatalytic reactions and spread the chain of damage by reacting with molecules and converting them into free radicals. Free radicals are mainly produced from these sources in the body: ubiquinone in the mitochondrial membrane, xanthine oxidase of endothelial cells, and myeloperoxidase and NADPH oxidase of neutrophils. But, xanthine oxidase and the respiratory chain of mitochondria are the major sources of reactive oxygen species (ROS) (Akhlaghi and Bandy, 2009). Free radicals are extremely unstable and become stable by pairing an outer shell electron with biomolecules, for example, lipids, proteins, and DNA. At high concentrations, free radicals can cause damage to various cell structures, comprising proteins and nucleic acids, together with lipid peroxidation (Poli *et al.*, 2004; Abraham *et al.*, 1999). These injuries and damages are the major contributions towards aging, cardiovascular diseases, atherosclerosis, cancer, and inflammatory diseases (Hazra *et al.*, 2008). Living cells contain a defensive system of antioxidants against the ROS which avoids their unnecessary production and inactivate them. Various antioxidants have been reported to protect the body from the damaging effects of oxidative stress. Nowadays, research has been increased in the area of finding novel natural antioxidants due to low side effects in comparison to synthetic antioxidants (Gorinstein *et al.*, 2003).

Cyclophosphamide (CYP), Figure 1a, is a synthetic alkylating agent chemically

related to the nitrogen mustards (Takimoto *et al.*, 2005) widely used as an anticancer and immunosuppressive drug (Clern and Bickers, 1991; Paul and Bruce, 1991) and in the treatment of nephrotic syndrome (Kirkland *et al.*, 1976; Etteldorf *et al.*, 1976). It is effective against a wide spectrum of malignancies, such as leukemia, lymphoma, breast, lung, prostate, and ovarian cancers (Khan *et al.*, 2004; Shanafelt *et al.*, 2007). Many anticancer drugs are known for the generation of Reactive Oxygen Species (ROS) in cancer cells (Hanane *et al.*, 2012) and these ROS generated lead to oxidative damage in the cell (Maiti, 2012). During the bioactivation of CYP, reactive oxygen species are also formed, which can modify the components of both healthy and neoplastic cells leading to decreased antioxidative capacity (Stankiewicz *et al.*, 2002). CYP's antineoplastic effects are associated with phosphoramidate mustard, while acrolein is linked with its toxic side effects (Kern and Kehrer, 2002). The cytotoxic effects of CYP and other chemotherapeutic drugs result in part from their interaction with DNA leading to defective DNA, abnormal cell function, and cell death (Lee and Schmitt, 2009). Several studies suggest that antioxidant supplementation can influence the response to chemotherapy as well as the development of adverse side effects that result from treatment with antineoplastic agents (Weiji *et al.*, 1997).

The best source to get a range of novel herbal drugs is medicinal plants. *Moringa oleifera* is called "Miracle Vegetable" because it is both a medicinal and functional food (Verma *et al.*, 1976).

Moringa oleifera has the highest proportion of essential amino acids and significant quantities of minerals (Sabate, 2003) when analyzed. *Moringa oleifera* is rich in compounds like glucosinolates and isothiocyanates (Fahey *et al.*, 2001). Flowers contain pigments such as alkaloids, kaempferol, rhamnetin, isoquercitrin, and kaempferitin (Faizi *et al.*, 1994). Phytochemical screening is of paramount importance in identifying a new source of a therapeutically and industrially valuable compound having medicinal significance, to make the best and judicious use of available natural wealth. Consequently, the present study aimed to investigate the antioxidant and chemoprotective effect of MLE against CYP-induced oxidative stress in rats.

2. Materials and Methods

2.1. Chemicals

Cyclophosphamide, reduced glutathione (GSH), bovine serum albumin (BSA), All reagents were of the highest quality. Reduced glutathione, 1-chloro-2, 4-dinitrobenzene, thiobarbituric acid (TBA), potassium iodide, sodium hydroxide, and trichloroacetic acid (TCA), and all other chemicals were purchased from Sigma Chemical Company (Saint Louis, USA).

2.2. Preparation of Extract

Moringa oleifera leaf powder (were supplied by Ancient Greenfields Ltd, Coimbatore-641108, INDIA). The aqueous extract was prepared as described by Thilza *et al.* (2010). 100 g of ground plant material (*Moringa* leaf) was macerated in 1.5 L of boiled distilled water for one hour. The mixture was filtered through Whatman filter paper No. 3 and the filtrate obtained was evaporated to dryness using a rotator evaporator at 45°C. The extract obtained

(22, 9% yield) was stored at 4°C. The extract solution was prepared in distilled water each time before experimentation.

2.3. Experimental Design

Male Sprague Dawley rats (160–170 g) of seven weeks old were used as an animal model in this study. They were maintained in cages at room temperature of 23 ± 2 °C with a 12 h light/dark cycle and free access to water and feed. Rats were acclimatized to the laboratory environment for two weeks before the start of the experiments. The European Community Directive (86/609/EEC) and National rules on animal care have been followed. All procedures with animals were conducted strictly following guidelines approved by the Institute Animal Ethical Committee. Permission to handling with laboratory animals was obtained from the Institute's animal committee. During the experiments, maximum care was taken to minimize animal suffering and, also, the number of rats used was kept at a minimum. Twenty male rats were randomly distributed into 4 groups (5 rats/group):

- Group I (control): Rats received vehicle used for MLE and CYP for 14 days.
- Group II (*Moringa* leaf extract (MLE)): Rats were administered orally with MLE 200 mg/kg/day (Aja *et al.*, 2015) was administered orally through an intragastric tube every day and continued until rats were sacrificed) for 14 days.
- Group III (cyclophosphamide (CYP): CYP at a dose of 150 mg/kg (Kim *et al.*, 2013) was given through i.p. to rats.
- Group IV (MLE + CYP): received MLE for 7 consecutive days, before and after CYP injection at a dose of 150 mg/kg was injected once on day 7, 1 h after MLE administration

2.4. Animal Dissection

After the last treatment, rats were unfed for 24 h. Diethyl ether was used to anesthetize animals and then dissected the animals from the ventral side of the body. Blood was collected by piercing heart. Blood was collected in two types of tubes; that is, for serum analysis, it was stored in small falcon tubes which were then centrifuged to obtain serum; the rest of the blood was collected in EDTA containing tubes for whole blood analysis. From the dissected animal, livers were removed and placed in saline solution. The liver was stored in liquid Nitrogen at -70°C for antioxidant enzymes and tissue stress marker examination.

2.5. Serum Analysis

After dissection of animals, blood was collected in falcon tubes and centrifuged at 4000 rpm for 20 min at 4°C , after 30 minutes, to collect the serum samples. Serum biomarkers were determined using a commercial kit following manufacturers' instructions using a spectrophotometer (Shimadzu UV-VIS Recording 2401 PC, Japan). Serum samples were analyzed for total protein by the Biuret method according to Armstrong and Carr (1964). Albumin concentration was determined by the method of Doumas *et al.* (1977). Globulin concentration was determined as the difference between total protein and albumin. Serum alanine aminotransferase (ALT; EC 2.6.1.2) and aspartate aminotransferase (AST; EC 2.6.1.1) activities were determined using commercial kits obtained from Biodiagnostic kit (Cairo, Egypt). The principal reaction of the colorimetric determination of AST or ALT activity is based on the reaction of aspartate or alanine with α ketoglutarate to form oxaloacetate or pyruvate hydrazone formed with 2, 4-dinitrophenylhydrazine (Reitman and

Frankel, 1957). Serum alkaline phosphatase (ALP; EC 3.1.3.1) activity was measured at 405 nm by the formation of para-nitrophenol from para-nitrophenyl phosphate as a substrate (Rosalki and Foo, 1993). Serum lactate dehydrogenase (LDH; EC 1.1.1.27) was determined according to the method of Friedman and Young (1997), using a kit obtained from Spinreact (Santa Coloma, Spain). Also, low-density lipoprotein (LDL) cholesterol was measured according to the method of Assmann *et al.* (1984), high-density lipoprotein (HDL) cholesterol was measured according to the method of Burstein *et al.* (1984). Cholesterol and triglycerides were measured according to the method Carr *et al.* (1993) using a Biodiagnostic kit (Cairo, Egypt).

2.6. Lipid peroxidation, antioxidant non-enzymatic and enzymatic estimation in liver tissue

Tissue homogenate was prepared by homogenizing 100 mg of liver tissue in 1 mL of 100 mM potassium phosphate buffer containing 1 mM EDTA at pH 7.4. The supernatant was collected in clean falcon tubes after centrifugation for 30 min at $12000 \times g$ at 4°C and was used for further analysis. The following assays were carried out to analyze the pharmacological activity against the toxicity induced with CYP in rats.

Lipid peroxidation assay

Using TBA, thiobarbituric acid reactive substances (TBARS) were measured in the tissue homogenate (Ohkawa *et al.*, 1979). An aliquot of 100 μL of tissue homogenate was added to 100 μL of 100 mM ascorbic acid, 10 μL of 100 mM FeCl_3 , and 290 μL of sodium phosphate buffer (pH 7.4). The reaction solution was incubated for 1 hour in a shaking water bath at 37°C . The addition of 500 μL of 10% TCA stopped the reaction. Place reaction tubes were placed in a boiling water bath for 15 minutes after adding

500 μ L of 0.67% TBA. After 15 min tubes were shifted on crushed ice for 5 minutes and centrifuged for 10 minutes at 2500 \times g. The optical density of supernatant was measured at 535 nm to determine the amount of TBARS formed. Using a molar extinction coefficient of 1.560×10^5 M/cm, lipid peroxidation activity was measured as an amount of TBARS formed/min/mg tissue.

Reduced glutathione

Reduced glutathione (GSH) activity was measured according to the method of Jollow *et al.* (1974). 500 μ L of tissue homogenate was precipitated by the addition of 500 μ L of sulfosalicylic acid (4%). After 1 hour incubation at 4°C, samples were centrifuged for 20 min at 1200 \times g. 33 μ L of supernatant was collected and added to aliquots containing 900 μ L of 0.1 M potassium phosphate buffer (pH 7.4) and 66 μ L of 100 mM DTNB. Reduced glutathione reacts with DTNB and forms a yellow colored complex. Absorption was measured at 412 nm. μ M GSH/g tissue represents GSH activity

Catalase (CAT) activity

Measurement of CAT (EC. 1.11.1.6) activity is based on the methodology of Shah *et al.* (2013), which relies on the decomposition of H₂O₂. An aliquot of 25 μ L of tissue homogenate was added to 100 μ L of 10 mM H₂O₂ and 625 μ L of 5 mM EDTA buffer (pH 8.0). The disappearance of H₂O₂ in the reaction mixture by catalase was measured spectrophotometrically at 230 nm. CAT activity was expressed as U/g tissue.

2.7. Measuring RNA and DNA concentration

The RNA concentration in the samples was determined by precipitation of the nucleic acids in 0.5 M HClO₄, after which the RNA was hydrolyzed ice by incubation in 0.3 M KOH at 60°C for 1 h.

After removal of DNA by 0.5 M HClO₄ precipitation, the RNA concentration was determined by boiling the samples for 30 min in 6 M HCl, 0.01% FeCl₃, and 0.3% orcinol. Absorption was measured at 660 nm. RNA concentrations were calculated using a calibration curve with highly purified total adult heart RNA that was isolated by ultracentrifugation through a cesium chloride cushion (Chirgwin *et al.*, 1979) as a standard.

The DNA concentration in the samples was determined by hydrolyzing the RNA in 0.1 M NaOH after which the DNA was precipitated with half volume 10% HClO₄. The DNA was suspended in 10% 32PHClO₄ and incubated at 70°C to hydrolyze the DNA. After clearing the solution by centrifugation, diphenylamine and acetaldehyde were added to a final concentration of 2% and 0.01%, respectively. After overnight incubation at 30°C, the absorbance was determined at 560 and 700 nm. The concentration in the samples was calculated relative to the included calibration curve that was prepared using highly Mopurified herring testis DNA (Sambrook *et al.*, 1989).

2.8. Statistical Analysis

Data are expressed as mean \pm S.E.M. (Significance was calculated at P<0.05). Statistical analysis was carried out using one-way analysis of variance (ANOVA) followed by Tukey-Kramer multiple comparisons test (Daniel, 1991).

3. Results

The present experiment was carried out to determine the protective role of MLE against CYP-induced liver toxicity at a biochemical level in rats.

3.1. Estimation of cyclophosphamide toxicity on serum protein, albumin, and globulin

Cyclophosphamide treatment in rats caused a significant change in the serum protein profile. To analyze the protective effect of MLE, the fluctuations in serum total protein, albumin, and globulin were analyzed (Table 1). A significant difference was observed in the CYP treated group in comparison to the control group. MLE treatment showed a protective role by significantly ($P < 0.05$) decreasing its level in the serum.

3.2. Hepatic toxicity

Liver function tests commonly include tests to measure levels of several enzymes, which are special proteins that help the body break down and metabolize other substances.

Enzymes that are often measured include ALT; AST and ALP (Berk and Korenblat, 2007). The protective effect of MLE on CYP – induced changes in serum ALT, AST, and ALP in rats was presented in Table (1). Treatment with CYP significantly ($P < 0.05$) increased the activities of AST, ALT, ALP, and LDH in serum compared to control animals. The present study demonstrates that treatment with MLE alone did not cause any significant change in enzyme activities in serum and alleviated the toxicity of CYP. MLE in combination with CYP alleviated its negative effect on activities of the above-measured enzymes (Table 1).

Table 1 Effects of Moringa leaf extract on biochemical parameters of serum from Cyclophosphamide-treated rats

Parameters	Groups			
	Control	MLE	CYP	MLE + CYP
Total protein (g/dl)	9.64 ± 0.69	9.88 ± 0.53	7.2 ± 0.97 ^{abd}	8.62 ± 0.25 ^{abc}
Albumin (g/dl)	4.38 ± 0.09	4.47 ± 0.13	4.13 ± 0.1 ^{abd}	4.41 ± 0.06 ^c
Globulin (g/dl)	5.86 ± 0.37	5.88 ± 0.24	3.3 ± 0.26 ^{abd}	4.78 ± 0.34 ^{abc}
AST (U/L)	122.2 ± 8.2	113.6 ± 2.72	310.0 ± 16.44 ^{abd}	209.6 ± 17.17 ^{abc}
ALT (U/L)	31.8 ± 1.7	27.4 ± 2.05	120.6 ± 013.06 ^{abd}	61.8 ± 17.08 ^{abc}
ALP (U/L)	154.0 ± 12.0	129.2 ± 12.9 ^{acd}	309.0 ± 24.5 ^{abd}	215.6 ± 19.9 ^{abc}
LDH (U/L)	1096.8 ± 81.4	918.6 ± 47.3 ^{acd}	1722.2 ± 152.9 ^{abd}	1378.2 ± 67.3 ^{abc}

Values are mean ± SE of five rats in each group. Significance at $P < 0.05$. MLE: Moringa leaf extract; CYP: Cyclophosphamide

^a Compared of control and other groups

^b Compared of MLE group and other groups

^c Compared of CYP group and other groups

^d Compared of MLE+CYP group and other groups

3.3. Lipid and lipoprotein profiles

Data represented in Table 2 show that i.p. treatment with CYP caused a significant ($P < 0.05$) increase in serum cholesterol, triglycerides, and LDL concentrations, whilst HDL levels were decreased compared to control animals. MLE alone caused a significant ($P < 0.05$) decrease in lipid profile compared to control. MLE-pretreated rats treated with CYP showed a reduction in the serum lipid level caused by CYP (Table 2).

Table 2: Effects of Moringa leaf extract on lipid and lipoprotein profiles of serum from Cyclophosphamide-treated rats

Parameters (mg/dl)	Groups			
	Control	MLE	CYP	MLE + CYP
Triglycerides	65.9 ± 3.08	52.48 ± 3.84 ^{acd}	111.08 ± 13.16 ^{abd}	87.68 ± 3.8 ^{abc}
Total cholesterol	50.98 ± 3.98	42.26 ± 1.8 ^{acd}	87.16 ± 6.08 ^{abd}	63.68 ± 5.85 ^{abc}
HDL	33.05 ± 2.1	37.08 ± 1.28 ^{acd}	20.86 ± 2.81 ^{abd}	27.61 ± 1.06 ^{abc}
LDL	14.42 ± 0.78	11.72 ± 1.0 ^{acd}	35.76 ± 5.09 ^{abd}	20.8 ± 2.84 ^{abc}

Values are mean ± SE of five rats in each group. Significance at $P < 0.05$. MLE: Moringa leaf extract; CYP: Cyclophosphamide; HDL, high-density lipoprotein cholesterol; LDL, low-density lipoprotein cholesterol.

^a Compared of control and other groups

^b Compared of MLE group and other groups

^c Compared of CYP group and other groups

^d Compared of MLE+CYP group and other groups

3.4. Lipid peroxidation, antioxidant non-enzymatic and enzymatic estimation in liver tissue

To characterize the protective effect of MLE, change in oxidant and antioxidant enzyme level was evaluated after CYP treatment. Figures 1-3 show the protective effect of MLE on liver tissue TBARS, GSH, and CAT. MLE per se did not affect serum levels of TBARS when compared with controls. In CYP-treated rats, a significant ($P < 0.001$) increase of lipid peroxidation in serum was observed, as evidenced by the enhanced TBARS levels in serum compared with those of controls. With MLE pretreatment in CYP-treated rats, a dramatic reduction ($P < 0.05$) in TBARS levels was recorded compared to the CYP-treated group (Figure 1).

In comparison to the control group, the levels of GSH and CAT in liver tissues were considerably decreased ($P < 0.05$) after the CYP treatment. MLE co-treatment reversed the activity of these non-enzymatic and enzymatic antioxidants towards normal (Figures 2, 3).

3.5. Liver DNA and RNA concentration

Cyclophosphamide decreased ($P < 0.05$) liver DNA and RNA concentrations compared to the control. The combination of MLE and CYP was less effective in DNA and RNA compared to individual treatment of CYP (Figure 4).

4. Discussion

Regardless of the enormous advancement in the field of pharmacology and conventional chemistry in creating effective medicines, still plant kingdom is a reservoir of natural therapeutics and offers a valuable source of novel drugs and medicinal entities. Medicinal plants and their phytochemicals are the main sources of herbal drugs that can affect the physiological system of animals

either directly or indirectly. Plant-based medicines have minimal or no side effects; therefore, these medicines are acknowledged for the treatment of some diseases. Various plants derived antioxidant-based therapeutic medicines are being in use for the prevention and cure of many diseases such as Alzheimer's disease, diabetes, stroke, atherosclerosis, and cancer (Devasagayam *et al.*, 2004). The current study mainly concentrates on the role of MLE in ameliorating the CYP induced oxidative stress in rats. It is evident from the results that MLE ethanol extract possess protective action against liver injuries induced by CYP induced oxidative stress.

A decrease in the level of serum protein, albumin, and globulin was observed after the CYP treatment in rats. The reduction in serum protein, particularly albumin, could be attributed to changes in protein and free amino acid metabolism and their synthesis in the liver (Attia and Nasr, 2009). Also, the protein level suppression may be due to loss of protein either by the reduction in protein synthesis or increased proteolytic activity or degradation (Yeragi *et al.*, 2003). Also, the observed decrease in serum proteins could be attributed in part to the damaging effect of CYP on liver cells, as confirmed by the increase in activities of serum AST and ALT. Natural antioxidant (MLE) mitigated the changes in the protein by regulating metabolic activities and protein synthesis.

In the current study, CYP treatment greatly affected hematological parameters. Serum ALT, AST, and ALP are sensitive markers employed in the diagnosis of liver damage (Boone *et al.*, 2005; Singh *et al.*, 2011). The increased activity of plasma ALP, ALT, AST, and level of LDH in the CYP-treated rats is a manifestation of induced hepatocellular damage. Increases in the ALP are generally associated with impairment of intrahepatic and extrahepatic bile flow,

hepatobiliary injury, erythrocyte destruction, or altered bilirubin metabolism (Singh *et al.*, 2011; Dufour *et al.*, 2000; Burt and Day, 2002). Chemotherapy-induced increase in plasma ALT and AST has been well documented (Ramadori and Cameron, 2010). Increases in ALT and AST activities are indicative of hepatocellular injury or regenerative activity (Meyer and Harvey, 2004). MLE could have a protector effect against hepatic damage induced by oxidative stress in the CYP state, which was evidenced by the capacity of this polyphenol to modulate the antioxidant defense and to decrease the lipid peroxidation in the liver tissues (Schmtz *et al.*, 2012).

Results also showed a significant increase in serum HDL of animals pretreated with MLE before CYP treatment compared to the CYP-treated group. HDL may hasten the removal of cholesterol from peripheral tissue to the liver for catabolism and excretion. Also, a high level of HDL may compete with LDL receptor sites on arterial smooth muscle cells and thus partially inhibit uptake and degradation of LDL. Also, HDL could protect LDL against oxidation *in vivo*, because the lipids in HDL are preferentially oxidized before those in LDL (Bowry *et al.*, 1992). These alterations in the lipid profile may be attributed to the increased lipolysis and fatty acid formation in the blood. Hypertriglyceridemia in combination with abnormally low concentrations of HDL is one of the most common atherogenic profiles of lipid metabolism. Agbor *et al.* (2005) reported that the accumulation of blood triglycerides may be a result of an imbalance between the rate of synthesis and the rate of release of triglycerides by the parenchyma cells into the systemic circulation.

Various studies have confirmed that free radicals are involved in various metabolic alterations and diseases. Radical reactions are mainly responsible for the *in*

vivo toxic effects of CYP. Lipids confined in the cell membranes are very sensitive to oxidative stress. Lipid peroxidation converts polyunsaturated fatty acids into small and more reactive elements. CYP is a toxic chemical that produces various free radicals causing lipid peroxidation. An increase in lipid peroxidation is calculated in terms of thiobarbituric acid reacting substance (TBARS) which measures the damage caused to the membranes by free radicals (Attia and Nasr, 2009). So, TBARS level content is supposed to be the main markers of CYP-induced oxidative stress (Stankiewicz *et al.*, 2002). In the current study, a decrease in protein content was observed while TBARS level increased due to CYP treatment which was restored to normal level after treating with MLE. During the bioactivation of CYP, reactive oxygen species are also formed, which can modify the components of both healthy and neoplastic cells leading to decreased antioxidative capacity (Stankiewicz *et al.*, 2002). Free radicals produced by bioactivation of CYP increase lipid peroxidation and decrease the level of antioxidant enzymes resulting in damage to liver tissue. MLE restored the damage produced by CYP. In line with the present observation Chattopadhyay *et al.* (2011), found that administration of aqueous extract of *M. oleifera* was found to significantly prevent the arsenic-induced alteration of hepatic function markers and lipid profile.

Various antioxidant enzymes, for example, catalase, peroxidase, and superoxide dismutase play an important role in protecting lung tissues from free radical-induced damages (Ganie *et al.*, 2011). In the present study, the decrease in the level of antioxidant enzymes (CAT) and GSH in liver tissues were observed in CYP treated group. Co-administration of MLE ameliorated the level of antioxidant enzymes, suggesting the protective role of antioxidant enzymes against CYP generated free radicals. Similar results were observed by Rakesh and Singh (2010) *M. oleifera* extract restored GSH in rat liver reduced by CCl₄ treatment.

The maintenance of GSH activity in a cell is dependent on the level of glutathione reductase and NADH (Meister and Anderson, 1983). A significant decrease in activities of GSH was observed in the present study. The decreased activity of the glutathione system in liver tissue of CYP intoxicated rats might be due to the increased lipid peroxidation or inactivation of the antioxidative enzymes. Administration of MLE ameliorated the CYP toxicity, thereby increasing the activity of antioxidant enzymes and GSH. Thus, activation of these antioxidants by the administration of MLE clearly shows that its free radical scavenging activity could exert a beneficial action against pathophysiological alterations caused by free radicals.




Figure 1: Liver thiobarbituric acid reactive substances (TBARS) (U/g tissue) of rats treated with *Moringa oleifera* leaf extract and cyclophosphamide. MLE: *Moringa oleifera* leaf extract; CYP: Cyclophosphamide; Significance at $P < 0.05$. ^a Comparison of control and other groups; ^b Comparison of MLE and other groups; ^c Comparison of CYP and other groups; ^d Comparison of MLE-CYP and other groups

Under normal conditions to protect cells against oxidative damage, the potentially harmful effects of ROS and free radicals are effectively eliminated by the antioxidant defense systems such as antioxidant enzymes and nonenzymatic factors (Hu *et al.*, 2005). Normally, cells possess a well-developed biochemical defense system, comprising low-molecular-weight free radical scavengers, i.e. glutathione (GSH), vitamin C, vitamin E, and complex enzymes, including GPx, SOD, and CAT (Yao *et al.*, 2007). When the balance between the generation and elimination of ROS is broken, as a result of these events, biomacromolecules including DNA, membrane lipids, and proteins are

damaged by ROS-mediated oxidative stress (Qian *et al.*, 2008; Ding and Ma, 2018).

Cyclophosphamide decreased liver DNA and RNA concentrations compared to the control. Supporting the present study, Matalon *et al.* (2004) reported that CYP requires microsomal oxidation in the liver to yield its active metabolite, 4-hydroxy-CYP in target cells that spontaneously decomposes to phosphoramidate mustard which exerts cytotoxic effect by the induction of DNA single-strand breaks as well as crosslinks which result in different types of damage. Also, Bosanquet *et al.* (2002) had reported that CYP involves inhibition of cell division due to cross-linking of the drug to DNA.




Figure 2: Liver glutathione (U/g tissue) of rat treated with *Moringa oleifera* leaf extract and cyclophosphamide. MLE: *Moringa oleifera* leaf extract; CYP: Cyclophosphamide; Significance at $P < 0.05$. ^a Comparison of control and other groups; ^b Comparison of MLE and other groups; ^c Comparison of CYP and other groups; ^d Comparison of MLE-CYP and other groups




Figure 3: Liver catalase activity (U/g tissue) of rat treated with *Moringa oleifera* leaf extract and cyclophosphamide. MLE: *Moringa oleifera* leaf extract; CYP: Cyclophosphamide; Significance at $P < 0.05$. ^a Comparison of control and other groups; ^b Comparison of MLE and other groups; ^c Comparison of CYP and other groups; ^d Comparison of MLE-CYP and other groups




Figure 4: Liver DNA and RNA concentrations (µg/gm tissue) of rats treated with *Moringa oleifera* leaf extract and cyclophosphamide. MLE: *Moringa oleifera* leaf extract; CYP: Cyclophosphamide; Significance at $P < 0.05$. ^a Comparison of control and other groups; ^b Comparison of MLE and other groups; ^c Comparison of CYP and other groups; ^d Comparison of MLE-CYP and other groups

The combination of CYP and MLE was less effective for DNA and RNA compared to the individual administration of CYP. In line with the present observation, Ippoushi *et al.*, (2003) found protection against oxidative DNA damage by herbal extracts and formulations. Oxidative damage mediated as single-strand breaks in supercoiled PTZ18U plasmid DNA has been reported to be suppressed by 6-gingerol (a phenolic compound in ginger). Also, a reduction in the basal levels of oxidative DNA damage upon treatment with 4-coumaric and protocatechuic acids was reported (Guglielmi *et al.*, 2003). The results of the present study are also in agreement with the above reports. These findings support the use of the leaves of *Moringa oleifera* to protect against oxidative DNA damage.

5. Conclusions

It can be concluded that *Moringa* extract contains antioxidant activity as it prevented oxidative stress and increased the antioxidant effect in liver tissues of male rats. Our results demonstrate the CYP protective role against CYP generated free radicals damages and suggestions for further study to isolate the bioactive component in pure form from MLE.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

Abraham, P., Wilfred, G., Cathrine, S.P. (1999). Oxidative damage to the lipids and proteins of the lungs, testis, and kidney of rats during carbon tetrachloride intoxication, *Clinica Chimica Acta*, 289(1-2): 177–179.

Agbor, G.A., Oben, J.E., Nkegoum, B., Takla, J.P., Ngogang, J.Y. (2005).

Hepatoprotective activity of *Hibiscus cannabinus* (Linn.) against carbon tetrachloride and paracetamol induced liver damage in rats. *Pak. J. Biol. Sci.*, 8: 1397–1401.

Aja, P.M., Ibiam, U.A., Nwali, B.U., Orji, O.U., Edwin, N., Afiukwa, C.A. (2015). Evaluation of Effect of Ethanol Extracts of *Moringa oleifera* and *Cajanus cajan* Leaves on Lipid Profiles in Alloxan Induced Diabetic Albino Rats. *GJBB*. 10 (2): 71-76.

Akhlaghi, M., Bandy, B. (2009). Mechanisms of flavonoid protection against myocardial ischemia-reperfusion injury. *J. Mol. Cell. Cardiol.*, 46(3): 309–317.

Armstrong, W.D., Carr, C.W. (1964). Estimation of serum total protein. In: *Physiological Chemistry Laboratory direction*, third ed. Minneapolis, Minnesota, USA: Burges Publishing Co.

Assmann, G., Jabs, H.U., Kohnert, U., Nolte, W., Schriewer, H. (1984). LDL-cholesterol determination in blood serum following precipitation of LDL with polyvinylsulfate. *Clin. Chem. Acta.*, 140: 77-83.

Attia, A.M., Nasr, H.M. (2009). Dimethoate-induced changes in biochemical parameters of experimental rat serum and its neutralization by black seed (*Nigella sativa* L.) oil. *Slovak J. Anim. Sci.*, 42(2): 87–94

Bosanquet, A.G., Sturm, I., Weider, T., Essmann, F., Bosanquet, M. I., Head, D. J., Dorken, B., Daniel, P. T. (2002): Bax expression correlates with cellular drug sensitivity to doxorubicin, cyclophosphamide and chlorambucil but not to fludrabin, cladribin or cortico-steroids in B cell chronic lymphocytic leukemia. *Leukemia J.*, 16: 1035.

Bowry, V.W., Stanley, K.K., Stocker, R. (1992). High density lipoprotein is the major of lipid hydroperoxides in human blood

- plasma from fasting donors. Proc. Natl. Acad. Sci., 89: 1316-1320
- Burstein, M., Scholnick, H.R., Morfin, R. (1980). Rapid method for the isolation of lipoproteins from human serum by precipitation with polyanions. Scand. J. Clin. Lab. Invest., 40: 583-595.
- Burt, A.D., Day, C.P. (2002). Pathophysiology of the liver In R.M.N. MacSween, A.D. Burt, B.C. Portmann, K.G. Ishak, P.J. Scheuer, P.P. Anthony eds Pathology of the Liver 4th ed London UK Churchill Livingstone. Pp 67–106.
- Carr, T., Andressen, C.J., Rudel, L.L. (1993). Enzymatic determination of triglyceride, free cholesterol and total cholesterol in tissue lipid extracts. Clin. Chem., 26: 39-42.
- Chattopadhyay, S., Maiti, S., Maji, G., Deb, B., Pan, B., Ghosh, D. (2011). Role of Protective Moringa oleifera (Sajina) seed on arsenic-induced hepatocellular degeneration in female albino rats. Biol. Trace Elem Res., 142(2): 200-212.
- Chirgwin, J.M., Przybyla, A.E., Macdonald, J., Rutter, W.J. (1979). Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. Biochemistry 18: 5294–5299
- Clern, M., Bickers, D.R. (1991). Dermatological pharmacology. In The Pharmacological Basis of Therapeutics; Goodman, G.A., Rall, T.W., Nies, A.S., Taylor, P., Eds.; Pergamon Press: New York, NY, USA, pp. 1572–1591.
- Daniel, W.W. (1991). Hypothesis testing. In: Biostatistics: A Foundation for Analysis in the Health Sciences: 5th ed. Jhon Wley & Sons. New York, Chichester, Brisbane, Toronto and Singapore.
- Devasagayam, T.P.A., Tilak, J.C., Bloor, K.K., Sane, K.S., Ghaskadbi, S.S., Lele, R.D. (2004). Free radicals and antioxidants in human health: Current status and future prospects. J. Assoc. Physicians India, 52: 794–804.
- Ding, H.M., Ma, T.Q. (2018). Computational approaches to cell–nanomaterial interactions: keeping balance between therapeutic efficiency and cytotoxicity. Nanoscale Horiz., 3: 6-27.
- Doumas, B.T., Watson, W.A., Biggs, H.G. (1977). Albumin standards and the measurement of serum albumin with bromocresol green. Clin. Chem. Acta., 31: 87-96
- Dufour, D.R., Lott, J.A., Nolte, F.S., Gretch, D.R., Koff, R.S., Seeff, L.B. (2000). Diagnosis and monitoring of hepatic injury II recommendations for use of laboratory tests in screening diagnosis and monitoring. Clin. Chem., 46: 2050–2068.
- Etteldorf, J.N., West, C.D., Pitcock, J.A., Williams, D.L. (1976). Gonadal function, testicular histology, and meiosis following cyclophosphamide therapy in patients with nephrotic syndrome. J. Pediatr., 88: 206–212.
- Friedman, K., Young, D.S. (1997). Effects of disease on clinical laboratory tests, 3rd ed. AACC press
- Ganie, S.A., Haq, E., Hamid, A., Qurishi, Y., Mahmood, Z., Zargar, B.A., Masood, A., Zarga, M.A. (2011). Carbon tetrachloride induced kidney and lung tissue damages and antioxidant activities of the aqueous rhizome extract of Podophyllum hexandrum, BMC Complement. Alternat. Med., vol. 11, article 17, 2011.
- Gorinstein, S., Yamamoto, K., Katrich, E. et al., (2003). Antioxidative properties of Jaffa sweets and grapefruit and their influence on lipid metabolism and plasma antioxidative potential in rats. Biosci. Biotechnol. Biochem., 67(4): 907–910.

- Guglielmi, F., Luceri, C., Giovannelli, L., Dolaro, P., Lodovici, M. (2003) Effect of 4-coumaric and 3, 4 dihydroxy benzoic acid on oxidative DNA damage in rat colonic mucosa. *Brit J Nutr.*, 89:581–587.
- Hanane, A., Claire, L., Caroline, A., Fabrice, M. (2012). Anticancer Drug Metabolism: Chemotherapy Resistance and New Therapeutic Approaches. In *Topics on Drug Metabolism*; Paxton, J., Ed.; InTech: Rijeka, Croatia.
- Hazra, B., Biswas, S., Mandal, N. (2008). Antioxidant and free radical scavenging activity of *Spondias pinnata*, *BMC Complementary and Alternative Medicine*, 8(1): article 63.
- Hu, Y., Rosen, D.J., Zhou, Y. (2005). Mitochondrial manganese superoxide dismutase expression in ovarian cancer: role in cell proliferation and response to oxidative stress. *J. Biol. Chem.*, 280:39485–39492.
- Jollow, D.J., Mitchell, J.R., Zampaglione, N. and Gillette, J.R. (1974). Bromobenzene induced liver necrosis. Protective role of glutathione and evidence for 3,4 bromobenzene oxide as the hepatotoxic metabolite. *Pharmacology*, 11(3): 151–169.
- Kern, J.C., Kehrer, J.P. (2002). Acrolein-induced cell death. A caspase-influenced decision between apoptosis and oncosis/necrosis. *Chem Biol Interact.*, 139: 79–95.
- Khan, T.S., Sundin, A., Juhlin, C., Wilander, E., Oberg, K. and Eriksson, B. (2004). Vincristine, cisplatin, teniposide, and cyclophosphamide combination in the treatment of recurrent or metastatic adrenocortical cancer. *Med. Oncol.*, 21: 167–177.
- Kim, S.-H., Lee, I.-C., Beak, H.-S., Moon, C., Kim, S.-H., Kim, J.C. (2013). Protective effect of diallyl disulfide on cyclophosphamide-induced testicular toxicity in rats. *Lab. Anim. Res.*, 29(4):204-211.
- Kirkland, R.T., Bongiovanni, A.M., Cornfield, D., McCormic, J.B., Parks, J.S., Tenore, A. (1976). Gonadotropin responses to leutinizing releasing factor in boys treated with cyclophosphamide for nephrotic syndrome. *J. Pediatr.*, 89: 941–944.
- Maiti, A.K. (2012). Genetic determinants of oxidative stress-mediated sensitization of drug resistant cancer cells. *Int. J. Cancer* 130: 1–9.
- Matalon, S.T., Ornoy, A., Lishner, M. (2004): Review of the potential effects of three commonly used antineoplastic and immunosuppressive drugs (cyclophosphamide, azathioprine, doxorubicin on the embryo and placenta). *Reproductive Toxicol.*, 18: 219 – 230
- Meister, A., Anderson, M.E. (1983). Glutathione, *Annual Rev. Biochem.*, 52: 711–760
- Lee, S., Schmitt, C.A. (2003). Chemotherapy response and resistance. *Curr. Opin. Genet. Dev.*, 13: 63–69.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J. (1951). Protein measurement with the Folin phenol reagent, *J. Biol. Chem.*, 193(1): 265–275
- Ohkawa, H.; Ohishi, N., Yagi, K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.*, 95: 351-358
- Paul, C., Bruce, A.C. (1991). Antineoplastic agents. In *The Pharmacological Basis of Therapeutics*; Goodman, G.A., Rall, T.W., Nies, A.S., Taylor, P., Eds; Pergamon Press: New York, NY, USA, pp. 1209–1263.
- Poli, G., Leonarduzzi, G., Biasi, F., Chiarpotto, E. (2004). Oxidative stress and cell signaling, *Curr. Med. Chem.*, 11(9): 1163–1182.

- Rakesh, S., Singh, VJ. (2010). *In vivo* antioxidant activity of Moringa oleifera leaf and pod extracts against carbon tetra chloride induced liver damage in albino mice. J. Chem. Pharm. Res., 2(6): 275-283.
- Ramadori, G., Cameron, S. (2010). Effects of systemic chemotherapy on the liver. Ann. Hepatol., 9(2): 133-143.
- Reitman, S., Frankel, S. (1957). A colorimetric method for the determination of serum glutamic oxaloacetic acid and glutamic pyruvic transaminases, Am. J. Clin. Pathol., 28: 56-63.
- Rosalki, S.B., Foo, A.Y., Burlina, A. (1993). Multicenter evaluation of iso-ALP test kit for measurement of bone alkaline phosphatase activity in serum and plasma. Clin. Chem., 39: 648-652.
- Yao, P., Li, K., Song, F., Zhou, S., Sun, X., Zhang, X., Nussler, A.K., Liu, L. (2007). Heme oxygenase-1 upregulated by Ginkgo biloba extract: potential protection against ethanol-induced oxidative liver damage. Food Chem. Toxicol., 45:1333–1342.
- Sambrook, J., Fritsch, E.F., Maniatis, T. (1989). Molecular cloning: a laboratory manual, 2nd edn. New York: Cold Spring Harbor Laboratory
- Schmatz, R., Perreira, L.B., Stefanello, N., Mazzanti, C., Spanevello, R., Gutierrez, J., Bagatini, M., Martins, C.C., Abdalla, F.H., Serres, J.D.S., Zanini, D., Vieira, J.M., Cardoso, A.M., Schetinger, M.R., Morsch, V.M. (2012). Effects of resveratrol on biomarkers of oxidative stress and on the activity of delta aminolevulinic acid dehydratase in liver and kidney of streptozotocin-induced diabetic rats. Biochimie, 94(2): 374-383
- Shah, N.A., Khan, M.R., Ahmad, B., Noureen, F., Rashid, U., Khan, R.A. (2013). Investigation on flavonoid composition and anti-free radical potential of Sida cordata. BMC Complement Altern. Med., 13(1): article 276.
- Shanafelt, T.D., Lin, T., Geyer, S.M., Zent, C.S., Leung, N., Kabat, B., Bowen, D., Grever, M.R., Byrd, J.C., Kay, N.E. (2007). Pentostatin, cyclophosphamide, and rituximab regimen in older patients with chronic lymphocytic leukemia. Cancer 109: 2291–2298.
- Singh, K.C., Kaur, R., Marar, T. (2011). Ameliorative effect of Vitamin E on Chemotherapy Iduced Side Effects in Rat Liver. J. Phamacol. Toxicol., 10: 39-43.
- Stankiewicz, A., Skrzydlewska, E., Makiela, M. (2002). Effects of amifostine on liver oxidative stress caused by cyclophosphamide administration to rats. Drug Metabol. Drug Interact. 19: 67–82.
- Takimoto, C.H., Calvo, E. (2005). Principles of oncologic pharmacotherapy. In Cancer Management: A Multidisciplinary Approach, 9th ed.; Pazdur, R., Coia, L.R., Hoskins, W.J., Wagman, L.D., Eds.; PRR Inc.: Seattle, WA, USA, pp. 23–42.
- Thilza, B., Sanni, S., Zakari, A.I., Sanni, F.S., Muhammed, T., Musa, B.J. (2010). *In vitro* antimicrobial activity of water extract of Moringa oleifera leaf stalk on bacteria normally implicated in eye diseases,”Academia Arena, 2(6): 80–82.
- Qian, Z.J., Jung, W.K., Kim, S.K. (2008). Free radical scavenging activity of a novel antioxidant peptide purified from hydrolysate of bullfrog skin, Rana catesbeiana shaw. Bioresource Technol., 99:1690–1698.
- Ippoushi, K., Azuma, K., Ito, H., Horic, H., Higashio, H. (2003) [6]-gingerol inhibits nitric oxide synthesis in activated J774–1 mouse macrophages and prevents peroxy nitrite induced oxidation and nitration reactions. Life Sci., 73:3427–3437.

Weiji, N.I., Cleton, F.J., Osanto, S. (1997). Free radicals and antioxidants in chemotherapy-induced toxicity. *Cancer Treat. Rev.* 1997, 23, 209–240.

Yeragi, S.G., Rana, A.M., Koli, V.A. (2003). Effect of pesticides on protein metabolism of mudskipper *Boleophthalmus*

dussumieri. *J. Ecotoxicol. Environ. Monit.*, 13:211-4.

Yulug, E., Türedi, S., Karagüzel, E., Kutlu, O., MenteSe, A., Alver, A. (2014). The short term effects of resveratrol on ischemia-reperfusion injury in rat testis. *J. Pediatr. Surg.*, 49(3): 484–489.